

4-17-2019

# Analyzing Change of Direction and the Laterally Resisted Split Squat: Incorporating a Lateral Vector into the Single Leg Squat

Derek Maddy

*Boise State University*

Chance Cooley

*Boise State University*

Tyler Brown

*Boise State University*

Shawn Simonson

*Boise State University*

---



# Analyzing Change of Direction and Laterally Resisted Split Squat

Derek Maddy, Chance Cooley, Tyler Brown, Shawn Simonson



BOISE STATE UNIVERSITY

## Incorporating a Lateral Vector into the Single Leg Squat

### INTRODUCTION

**Agility** – The ability to change directions in response to a stimulus<sup>1</sup>

- Cognitive – How quickly the brain processes information in response to a stimulus (motor learning)
- Physical – How the body applies the information to change directions (change of direction: COD)

**Frequently tested in athletes to predict success**<sup>2-4</sup>

**COD Characteristics**<sup>2</sup>

- Unilateral
- Horizontal and Lateral Force Production
- Hamstring & Gluteal Muscle Activation

### IMPROVING COD MOVEMENT

Dos’Santos et al analyzed mechanical determinants for faster COD in athletes<sup>5</sup>

- Participants with faster ground contact time and higher horizontal propulsive forces have quicker COD<sup>5</sup>
- Slower participants have greater vertical impact forces and smaller horizontal propulsive forces during COD<sup>5</sup>

**Mixed Results for Improving COD via Strength Training**

- Spiteri et al correlation of strength to COD<sup>6</sup>
  - Jullien et al – No improvement via concentric squat training<sup>7</sup>
  - McBride et al – Improvement via eccentric training & motor learning effect<sup>8</sup>
  - Castillo-Rodriguez et al – Improvement unilateral CMJ but injuries occurred<sup>9</sup>
  - Speirs et al - Modified Single Leg Squat (MSLS) and Bilateral Squat (BS) resulted in similar improvement in COD<sup>10</sup>

### FORCE VECTOR THEORY

**Specificity of Strength Training** – Training that targets the muscles used during athletic skills is best for direct transferability<sup>11</sup>

- Training adaptations may be direction-specific and exercises should exhibit concentric and eccentric loads in the same anatomical plane as in the athletic movement<sup>12-14</sup>

- For COD, a transverse plane force is required for improved performance


**There currently is no resistance training movement that does this.**

- MSLS targets muscles used during COD and is performed in unilateral fashion; however, the MSLS is performed in an upright position and resisting a vertical downward force
- COD is not performed in an upright vertical position. This may reduce the transferability from the MSLS to COD

**The Laterally Resisted Split Squat (LRSS) is a modification of the MSLS that incorporates a lateral force vector**



**Figure 1:** An American football player performing a change in direction (B), comparing force production of the MSLS (A) and LRSS (C).<sup>15</sup>



**Figure 2:** Pilot data comparing the GRF vector between a 90° COD at the start of the propulsive phase (A) and the LRSS during the concentric phase of the lift (B).

### PURPOSE AND HYPOTHESIS

The purpose of this study is to compare COD with the LRSS, MSLS and the BS via kinetic measurements and muscle activation.

**Hypothesis I:** It is hypothesized that horizontal ground reaction force (HGRF) of the LRSS will be no different than COD, but will be significantly different than the BS.

- a) It is hypothesized that horizontal GRF of the LRSS will be significantly different than the MSLS.

**Hypothesis II:** It is hypothesized that the LRSS will require greater muscle activation in the bicep femoris, semimembranosus, gluteus maximus, and gluteus medius than the BS.

- a) It is hypothesized that the LRSS will require greater muscle activation in the bicep femoris, semimembranosus, gluteus maximus, and gluteus medius than the MSLS.

### METHODS

**Participants**

12 recreational college athletes participating in COD sports

- No previous lower limb injury over last 6 months
- No previous lower limb surgical procedures
- Provide written consent & health history form

**Measurements**

- LRSS, MSLS, BS - Indirect 1RM<sup>16</sup>
- Peak HGRF ( $HGRF = ((ML^2 + A - P^2)/BM))^{1/2}$ ;  $AMTI\ FP^{11}$ )
- Peak Muscle Activation (MVC; Delsys Trigno)<sup>17</sup>


**Protocol**

- 1) COD mechanic screening
- 2) Familiarization of LRSS
- 3) Indirect 1RM – LRSS, MSLS, BS
- 4) COD on Dominant Leg (GRF)
- 5) MVC
- 6) 3 Movements 70% 1RM

**Data Analysis**

SPSS 25 ( $\alpha=0.05$ )

- RM-ANOVA
  - HGRF of LRSS, MSLS, BS
  - EMG Peak Activation of LRSS, MSLS, BS



**Figure 3:** EMG placements of surface electrodes.<sup>17</sup>

### REFERENCES

- 1) Gleason, BH, Kramer, JB, Stone, MH. Agility training for American football. *Nat Strength Cond J*, 2016; 37(6): 65-71.
- 2) Brugheili, M, Cronin, J, Levin, G, Chaouachi, A. Understanding change of direction ability in sport: A review of resistance training studies. *Sports Med*, 2008; 38(12): 1045-1063.
- 3) Robbins, DW, Goodale, TL, Kuzmits, FE, Adams, AJ. Changes in the athletic profile of elite college American football players. *J Strength Cond Res*, 2013; 27(4): 861-874.
- 4) Sierer, SP, Battaglini, CL, Mihalik, JP, Shields, EW, Tomasini, NT. The National Football League Combine: Performance differences between drafted and nondrafted players entering the 2004 and 2005 drafts. *J Strength Cond Res*, 2008; 22(1): 6-12.
- 5) Dos Santos, T, Thomas, C, Jones, PA, Comfort, P. Mechanical determinants of faster change of direction speed performance in male athletes. *J Strength Cond Res*, 2017; 31(3): 696-705.
- 6) Spiteri, T, Cochrane, JL, Hart, NH, Hall, GG, Nimphius, S. Effect of strength on plant foot kinetics and kinematics during a change of direction task. *Eur J Sport Sci*, 2013; 13(6): 646-652.
- 7) Jullien, H, Bisch, C, Laroque, N, Manouvier, C, Carling, C, Amard, V. Does a short period of lower limb strength training improve performance in field-based tests of running and agility in young professional soccer players? *J Strength Cond Res*, 2008; 22(2): 404-411.
- 8) McBride, JM, Triplett-McBride, T, Davis, A, Newton, RU. The effect of heavy- vs. light load jump squats on the development of strength, power, and speed. *J Strength Cond Res*, 2002; 16(1): 75-82.
- 9) Castillo-Rodriguez, A, Fernandez-Garcia, JC, Chinchilla-Minguel, JL, Camero, EA. Relationship between muscular strength and sprints with changes of direction. *J Strength Cond Res*, 2012; 26(3): 725-732.
- 10) Speirs, DE, Bennett, M, Finn, CV, Turner, A. Unilateral vs. Bilateral squat training for strength sprints and agility in academy rugby players. *J Strength*
- 11) Young, WB. Transfer of strength and power training to sports performance. *Int J Sports Phys Per*, 2006; 1: 74-83 and Res. 2016; 30(2): 386-392.
- 12) Contreras, B, Vigotsky, AD, Schoenfeld, BJ, et al. Effects of a six-week hip thrust vs. front squat resistance training program on performance in adolescent males: A randomized controlled trial. *J Strength Cond Res*, 2017; 31(4): 999-1008.
- 13) Loturco, I, Contreras, B, Kobal, R, et al. Vertically and horizontally directed muscle power exercises: Relationships with top-level performance. *PLoS ONE*, 2018; 13(7).
- 14) Gonzalo-Skokol, O, Sanchez-Sabate, J, Izquierdo-Lupon, L, Saez de Villarreal, E. Influence of force-vector and force application plyometric training in young elite basketball players. *Eur J Sport Sci*, 2018; 1-10.
- 15) McCurdy, K, O'Keefe, E, Kutz, M, Langford, G, Ernest, J, Torres M. Comparison of lower extremity EMG between the 2-leg squat and modified single-leg squat in female athletes. *J Sport Rehab*, 2010; 19: 57-70.
- 16) Brzycki, M. Strength testing: Predicting a one-rep max from reps-to-fatigue. *J Phys Ed, Rec Dance*, 1993; 64(1): 88-90.
- 17) Konrad, P. The ABC of EMG: A practical introduction to kinesiological electromyography. Noraxon INC, USA, 2005.